

Kiehl-RapiDés-plus

Liquid rapid disinfectant

- ▲ Ready-to-use disinfectant
- ▲ Effective against gram-positive and gram-negative bacteria (incl. salmonella and listeria), yeasts, enveloped viruses (incl. HIV, HBV, HCV, SARS-CoV-2), adenoviruses, noroviruses and rotaviruses (limited spectrum virucidal activity)
- ▲ Tuberculocidal and mycobactericidal
- ▲ Short drying time
- ▲ Once dry, the surfaces can be immediately reused
- ▲ No need to rinse with water
- ▲ Free from aldehydes, fragrances and dyes
- ▲ The product has a clearance certificate for use in food areas



Area of application

For the disinfection of alcohol-resistant surfaces. Do not use on acrylic glass, surfaces sensitive to alcohol or paint.

Application

Please test on a small inconspicuous spot before first application. Remove heavy, oily, greasy and protein-based stains in advance. Screw on a sprayer (aerosol-free) or spray nozzle depending on the area of application.

Surface disinfection:

Fully cover surfaces, devices and objects with the undiluted product and keep moist during the reaction time. Leave to dry or wait for the product to take effect and then wipe clean with a dry disposable towel. Consumption: 7.0–50.0 ml/m².



Note

Kiehl accepts no liability or responsibility for damage caused as a result of incorrect use or application of the product. For professional use and private consumers. Use biocides safely. Always read the label and product information before use.
PT 02, PT 04

Ingredients

Water-soluble solvents.
pH value (concentrate): approx. 5.5

Active substances in 100 g:

62.0 g Ethanol (CAS: 64-17-5).

Classification

Classification (in concentrate): GHS02, GHS07, Danger. H225 Highly flammable liquid and vapour. H319 Causes serious eye irritation. P210 Keep away from heat, hot surfaces, sparks, open flames and other ignition sources. No smoking. P280 Wear eye protection. P305+P351+P338 IF IN EYES: Rinse cautiously with water for several minutes. Remove contact lenses, if present and easy to do. Continue rinsing. P403+P233 Store in a well-ventilated place. Keep container tightly closed. Contains: Ethyl Alcohol (INC)

Art. No.	Packaging unit (PU)	Numbers of PU per pallet
j350547	Carton of 6 x 750 ml pear shaped bottles with 1 sprayer	64
j3505k47	Carton of 6 x 750 ml pear shaped bottles with 6 sprayers	64
j350505	Carton of 2 x 5 L canisters	56

Kiehl-RapiDés-plus

Liquid rapid disinfectant

Tests and listings

Recommended dosage for surface disinfection in the healthcare sector

Test method	Spectrum of activity	Organic load	Dosage	Contact time
EN 16615 Surface disinfection with mechanical action	bactericidal / yeasticidal	low	undiluted	2 min
EN 16777 Quantitative surface test	limited spectrum virucidal activity			

Recommended dosage for surface disinfection in industrial and institutional areas and food industry

Test method	Spectrum of activity	Organic load	Dosage	Contact time
EN 13697 Surface disinfection without mechanical action	bactericidal / yeasticidal	low	undiluted	2 min
EN 16777 Quantitative surface test	limited spectrum virucidal activity			

Further tests

Test method	Spectrum of activity	Organic load	Dosage	Contact time
Medical sector				
EN 13727 Quantitative suspension test	bactericidal	low	undiluted	15 s
EN 13624 Quantitative suspension test	yeasticidal	low	undiluted	15 s
EN 17387 Surface disinfection without mechanical action	bactericidal / yeasticidal	low	undiluted	3 min
VAH-certified application recommendation for surface disinfection without mechanical action	bactericidal / yeasticidal	low	undiluted	5 min
EN 14348 Quantitative suspension test	mycobactericidal	low	undiluted	1 min
EN 17387 Quantitative surface test	mycobactericidal	low	undiluted	1 min
Industrial and institutional areas, food industry				
EN 1276 Quantitative suspension test	bactericidal	low	undiluted	1 min
EN 1650 Quantitative suspension test	yeasticidal	low	undiluted	1 min
EN 13697 Surface disinfection without mechanical action	bactericidal / yeasticidal	low	undiluted	1 min

See also www.desinfektionsmittelliste.de

Kiehl-RapiDés-plus

Liquid rapid disinfectant

Test method	Spectrum of activity	Organic load	Dosage	Contact time
Virucidal activity				
EN 14476 Quantitative suspension test	limited spectrum virucidal activity	low	undiluted	2 min
EN 16777 Quantitative surface test	limited spectrum virucidal activity	low	undiluted	2 min
EN 14476 Quantitative suspension test	Murine Norovirus (MNV)	low	undiluted	1 min
	Adenovirus			2 min
EN 16777 Quantitative surface test	Murine Norovirus (MNV)	low	undiluted	1 min
	Adenovirus			2 min

See also www.desinfektionsmittelliste.de