

ProMop-DES-K

Liquid disinfectant cleaner concentrate

- ▲ Especially developed for use in the ProMop system
- ▲ Effective against gram-positive and gram-negative bacteria (incl. MRSA and MRE), yeasts, and non-enveloped and enveloped viruses (virucidal effect incl. HIV, HBV, HCV, adeno and polio viruses)
- ▲ Included in the IHO List of Disinfectants
- ▲ Meets the requirements of EN 13727, EN 13624 and EN 16615
- ▲ Effective against noroviruses
- ▲ Effective against SARS-CoV-2
- ▲ Optimum cleaning performance
- ▲ Streak-free finish



Area of application

For the cleaning and disinfection of water-resistant floor, wall and work surfaces. Can also be used on coated floorings.

Application

Please test on a small inconspicuous spot before first application.

Preparing the cleaning textiles:

Prior to using the product, prepare the cleaning textiles for use in the ProMop washing process and store in boxes until needed.

Disinfecting cleaning:

Attach the canister to the pump connection for the preparation product. The dosage is set by the washing machine's pump control. See table for dosage. Remove cleaning cloths from the boxes and clean the surfaces. Fully cover surface with disinfection solution (approx. 15–20 ml/m²) and clean with a cleaning cloth or cleaning device. Consumption: no indication possible.

The amount required depends on the size of the washing machine, the water hardness and the chosen liquor ratio.



Note

Kiehl accepts no liability or responsibility for damage caused as a result of incorrect use or application of the product. Not a consumer product according to 1999/44/EC Art. 1! For professional use only! Use biocides safely. Always read the label and product information before use. PT 02

Ingredients (according to 648/2004/EC)

Nonionic surfactants 5–15%, biocides.
pH value (concentrate): approx. 5

pH value (ready-to-use solution): approx. 8

Active substances in 100 g:

11.0 g Glutaraldehyde (CAS: 111-30-8), 2.0 g Didecyldimethylammonium chloride (CAS: 7173-51-5).

Classification

Classification (in concentrate): GHS05, GHS07, GHS08, GHS09, Danger. H302+H332 Harmful if swallowed or if inhaled. H314 Causes severe skin burns and eye damage. H317 May cause an allergic skin reaction. H334 May cause allergy or asthma symptoms or breathing difficulties if inhaled. H335 May cause respiratory irritation. H400 Very toxic to aquatic life. P260 Do not breathe dust/spray. P273 Avoid release to the environment. P280 Wear protective gloves/protective clothing/eye protection/face protection. P305+P351+P338 IF IN EYES: Rinse cautiously with water for several minutes. Remove contact lenses, if present and easy to do. Continue rinsing. P308+P313 IF exposed or concerned: Get medical advice/attention. Contains: Glutaral / Didecyldimonium Chloride (INCI)

Art. No.	Packaging unit (PU)	Numbers of PU per pallet
j650610	10 L canister	60

ProMop-DES-K

Liquid disinfectant cleaner concentrate

Tests and listings

Recommended dosage for surface disinfection in the healthcare sector

Test method	Spectrum of activity	Organic load	Dosage		Contact time	Test microorganisms
EN 16615 / 4-field test VAH-certified application recommendation for surface disinfection with mechanical action	bactericidal yeastocidal	high	0.5%	50 ml/10 L	15 min	Staphylococcus aureus incl. MRSA, Enterococcus hirae, Escherichia coli, Pseudomonas aeruginosa Candida albicans
DVV/RKI Quantitative suspension test	virucidal activity against enveloped viruses	with and without				BVDV (surrogate for HCV), Vaccinia-virus

Further tests

Test method	Spectrum of activity	Organic load	Dosage		Contact time	Test microorganisms
Medical sector						
EN 13697 Surface disinfection without mechanical action	bactericidal yeastocidal	high	0.5%	50 ml/10 L	15 min	Staphylococcus aureus incl. MRSA, Enterococcus hirae, Pseudomonas aeruginosa Candida albicans
EN 13727 Quantitative suspension test	bactericidal	high	0.5%	50 ml/10 L	5 min	Staphylococcus aureus incl. MRSA, Enterococcus hirae, Pseudomonas aeruginosa
EN 13624 Quantitative suspension test	yeastocidal	high	0.5%	50 ml/10 L	5 min	Candida albicans
Virucidal activity						
DVV/RKI Quantitative suspension test	virucidal	with and without	0.5%	50 ml/10 L	90 min	Enveloped and non-enveloped viruses
DVV/RKI Quantitative suspension test	virucidal activity against enveloped viruses	with and without	0.5%	50 ml/10 L	5 min	BVDV (surrogate for HCV), Vaccinia-virus
DVV/RKI Quantitative suspension test		with and without	0.5%	50 ml/10 L	30 min	Adenovirus Polyomavirus
DVV/RKI Quantitative suspension test		with and without	0.5%	50 ml/10 L	90 min	Poliovirus

See also www.desinfektionsmittelliste.de